



ORIGINAL ARTICLE

## *Azotobacter chroococcum* as a potentially useful bacterial biofertilizer for cotton (*Gossypium hirsutum*): Effect in reducing N fertilization



Felipe Romero-Perdomo<sup>a</sup>, Jorge Abril<sup>a</sup>, Mauricio Camelo<sup>a</sup>, Andrés Moreno-Galván<sup>a</sup>, Iván Pastrana<sup>a</sup>, Daniel Rojas-Tapias<sup>a,b</sup>, Ruth Bonilla<sup>a,\*</sup>

<sup>a</sup> Laboratorio de Microbiología de Suelos, Corporación Colombiana de Investigación Agropecuaria, Corpoica, Mosquera 250014, Colombia

<sup>b</sup> Graduate Field at Department of Microbiology, Cornell University, Wing Hall, Ithaca, NY 14853, USA

Received 17 November 2016; accepted 21 April 2017

Available online 31 August 2017

### KEYWORDS

Plant growth-promoting (rhizo)bacteria (PGPR/PGPB); Biofertilizer; Nitrogen-fixing bacteria; IAA; Phosphate solubilization

**Abstract** The aim of this research was to evaluate whether the application of two plant growth-promoting (rhizo)bacteria might reduce nitrogen fertilization doses in cotton. We used strains *Azotobacter chroococcum* AC1 and AC10 for their proven ability to promote seed germination and cotton growth. These microorganisms were characterized by their plant growth-promoting activities. Then, we conducted a glasshouse study to evaluate the plant growth promoting ability of these strains with reduced doses of urea fertilization in cotton. Results revealed that both strains are capable of fixing nitrogen, solubilizing phosphorus, synthesizing indole compounds and producing hydrolytic enzymes. After 12 weeks, the glasshouse experiment showed that cotton growth was positively influenced due to bacterial inoculation with respect to chemical fertilization. Notably, we observed that microbial inoculation further influenced plant biomass ( $p < 0.05$ ) than nitrogen content. Co-inoculation, interestingly, exhibited a greater beneficial effect on plant growth parameters compared to single inoculation. Moreover, similar results without significant statistical differences were observed among bacterial co-inoculation plus 50% urea and 100% fertilization. These findings suggest that co-inoculation of *A. chroococcum* strains allow to reduce nitrogen fertilization doses up to 50% on cotton growth. Our results showed that inoculation with AC1 and AC10 represents a viable alternative to improve cotton growth while decreasing the N fertilizer dose and allows to alleviate the environmental deterioration related to N pollution.

© 2017 Asociación Argentina de Microbiología. Published by Elsevier España, S.L.U. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

\* Corresponding author.

E-mail address: [rbonilla@corpoica.org.co](mailto:rbonilla@corpoica.org.co) (R. Bonilla).

**PALABRAS CLAVE**

(Rizo)bacteria promotora de crecimiento vegetal; Biofertilizante; Bacteria fijadora de nitrógeno; AIA; Solubilización de fosfato

### ***Azotobacter chroococcum* como biofertilizante bacteriano potencialmente útil para el algodón (*Gossypium hirsutum* L.): efecto en la reducción de la fertilización nitrogenada**

**Resumen** El objetivo de esta investigación fue evaluar si la aplicación de 2 (rizo)bacterias promotoras del crecimiento vegetal podría reducir la dosis de fertilizante nitrogenado en el cultivo de algodón. Se usaron las cepas *Azotobacter chroococcum* AC1 y AC10 por su habilidad para promover la germinación de semillas y el crecimiento del algodón. Estos microorganismos fueron caracterizados sobre la base de sus actividades de promoción del crecimiento vegetal. Luego se realizó un estudio de invernadero con plantas de algodón para evaluar la capacidad de promoción del crecimiento vegetal de dichas cepas con dosis reducidas de urea. Los resultados revelaron que ambas cepas son capaces de fijar nitrógeno, solubilizar fósforo, sintetizar compuestos indólicos y producir enzimas hidrolíticas. Después de 12 semanas, el experimento de invernadero permitió observar que el crecimiento del algodón fue influido positivamente por la inoculación bacteriana con respecto a la fertilización química. En particular, se evidenció que la inoculación microbiana impactó más en la biomasa vegetal ( $p < 0,05$ ) que en el contenido de nitrógeno. Curiosamente, la coinoculación exhibió un mayor efecto positivo sobre los parámetros de crecimiento en comparación con la inoculación simple. Además, se observaron resultados similares, sin diferencias estadísticamente significativas, entre la coinoculación bacteriana más del 50% de urea y el 100% de fertilización. Estos hallazgos indican que la coinoculación de las cepas de *A. chroococcum* AC1 y AC10 permitiría reducir las dosis de fertilización nitrogenada del cultivo de arroz en hasta el 50% y aliviar, de esta manera, el deterioro ambiental relacionado con la contaminación por N.

© 2017 Asociación Argentina de Microbiología. Publicado por Elsevier España, S.L.U. Este es un artículo Open Access bajo la licencia CC BY-NC-ND (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

## Introduction

Sustainable agricultural production requires new approaches to reduce the applications of polluting agrochemicals. In particular, the application of synthetic nitrogen fertilizers (SNF) represents an important environmental threat as it involves pollution due to its production and usage<sup>33</sup>. Within 40–50% of the total fertilizers involved in nitrogen uptake are not assimilated immediately by plants or are lost *via* leaching, denitrification, volatilization and are subject to conversion into unavailable forms. Poor assimilation of nutrients eventually leads to an increasing utilization of SNF, which reduces soil fertility, and biodiversity, contaminates groundwater, and consequently affects human health<sup>19</sup>. In Colombia, the unsustainable cotton production system is an example of the negative aftermath of SNF overuse. Low profitability for producers, lower export competitiveness, and environmental degradation are some consequences of this problem<sup>5</sup>. One potential solution to overcome this major challenge is the study of the biological processes involving plant growth-promoting (rhizo)bacteria (PGPR/PGPB) and their interaction with plants<sup>2</sup>.

The group of PGPR/PGPB encompasses several bacterial phyla, and although the direct molecular mechanisms associated to plant growth promotion remain poorly understood, it is widely accepted that their primary effect on plant growth is due to increasing availability of essential nutrients, plant growth stimulation, phytopathogen control, and decrease of negative effects in some abiotic stress

involved in plant growth<sup>8,9</sup>. Several lines of evidence suggesting direct mechanisms of PGPR/PGPB involved in plant growth promotion are the following: (i) production of ACC deaminase, which reduces the level of ethylene in crop roots thus enhancing root length and density; (ii) symbiotic and associative nitrogen fixation, which increases the availability of soluble nitrogen in soil; (iii) nutrient solubilization and mineralization (*e.g.* P, K, Zn and Si), which increases the availability of those elements for plant uptake; (iv) synthesis of phytohormones such as indoles, gibberellins, abscisic acid and cytokinins which modulate plant growth and division; (v) ability to produce siderophores, hydrolytic enzymes and antibiotics, which renders the cells more competitive over niche colonization; (vi) quorum sensing signal interference and inhibition of biofilm formation, and (vii) enhanced resistance to stresses by synthesis of water soluble vitamins as niacin, thiamine and biotin<sup>4,12</sup>. Undoubtedly, the relationship among those bacterial attributes are associated to plant growth promotion in diverse studies, and hence their use is promising as a biotechnological tool to be integrated into the agricultural production systems<sup>13</sup>.

Within the PGPR/PGPB group, the nitrogen-fixing bacterium *Azotobacter chroococcum* has shown to promote the growth of different crops under different soil types and climatic conditions<sup>15</sup>. Indeed, recent literature suggests that *A. chroococcum* might be a potential alternative to ameliorate the negative impact of NaCl on maize growth<sup>29</sup>. Although their positive effects on plant growth have been reported, few studies have explored the potential of *A. chroococcum* as supplement of chemical fertilization. For

example, Bonilla et al.<sup>5</sup> found that the application of *A. chroococcum* AC1 and AC10 increases cotton biomass. Moreover, significant differences in all plant growth parameters were evidenced between the inoculated treatment and the full fertilizer rate (positive control). They suggested that the use of AC1 and AC10 could not completely replace nitrogen (N) fertilizer. The goal of this work was to determine whether the inoculation of these *A. chroococcum* strains might reduce N fertilization rates in cotton. For this purpose, we evaluated the integrated use among these microorganisms and different doses of urea on plant growth attributes and nitrogen content under glasshouse conditions. Moreover, we related the evidenced results with the plant growth-promoting traits measured in these bacteria.

## Materials and methods

### Microorganisms and culture conditions

We used the *Gossypium hirsutum*-associated bacterial strains *A. chroococcum* AC1 and *A. chroococcum* AC10 for their compatibility, and ability to promote seed germination and growth of cotton without fertilization<sup>5</sup>. AC1 and AC10 strains were obtained from the Soil Microbiology Lab at Corpoica, Colombia. For routine use, both strains were stored at  $-80^{\circ}\text{C}$  in 50% (v/v) glycerol in MBR broth<sup>22</sup>, and grown on Ashby plates (composition in g/l: mannitol 10.0,  $\text{K}_2\text{HPO}_4$  0.2,  $\text{MgSO}_4$  0.2, NaCl 0.2,  $\text{CaSO}_4$  0.1,  $\text{CaCO}_3$  5.0, agar 15.0, pH 7.5) at standard conditions (*i.e.*  $30^{\circ}\text{C}$  for 48 h). All reagents used were from Merck, Millipore, USA.

### In vitro screening of plant growth promoting features

In order to characterize the plant growth promoting (PGP) traits of AC1 and AC10 strains, the ability to fix nitrogen<sup>11</sup>, solubilize nutrients as  $\text{P}^{10}$  and potassium<sup>17</sup>, produce the ACC deaminase enzyme<sup>16</sup>, synthesize Indole Acetic Acid (IAA)<sup>6,14</sup> and siderophores<sup>31</sup>, and secrete cell-wall-degrading enzymes (proteases and chitinases) and other enzymes such as pectinases, cellulases, amylases, and ureases<sup>7</sup> were evaluated. Each assay was performed with at least two biological replicates, and three technical replicates.

### Influence of *A. chroococcum* and fertilizers on cotton

In order to determine whether inoculation with AC1 and AC10 as single or mixed inoculation could reduce the fertilization dosage on cotton, we established a glasshouse experiment at Centro de Investigación Motilonia (CIM) of Corpoica, Agustín Codazzi, Colombia. The experiment was carried out in a completely randomized design (CRD) with six treatments. Each experiment was performed three times with six replicates. The fertilizer doses were 100, 75 and 50% (p/v) of urea (Ecofertil<sup>®</sup>, Colombia) with respect to the recommended application, each accompanied with 100% of both diammonium phosphate (DAP; Nutrimon<sup>®</sup>, Colombia) and KCl (Diabonos<sup>®</sup>, Colombia), as sources of N, P, K respectively; 100% of fertilizer corresponded to

75 kg/ha–2.34 g/pot for urea, 50 kg/ha–1.56 g/pot for DAP and 50 kg/ha–1.56 g/pot for KCl. We used *A. chroococcum* AC1, AC10, AC1+AC10 (1:1) strains, and sterile MBR broth as control treatment. In all cases, the biological treatments were applied along with 50% urea. Bacterial inoculum ( $\sim 1 \times 10^9$  CFU/ml) was prepared by growing the cells in MBR broth, using a Rushton type turbine-agitated fermenter (Miniforms, INF-30174) at 300 rpm, 1 lpm, and  $30^{\circ}\text{C}$  during 24 h. Soil was collected from cotton at CIM and its preparation was performed according to Rojas-Tapias et al.<sup>29</sup>. Cotton seedlings of DP ACALA 90 were surface-sterilized using 1% (v/v) NaClO for 10 min and rinsed three times in deionized-sterilized water. The surface-sterilized seeds were inoculated ( $\sim 4 \times 10^8$  CFU/ml/seed) by soaking with 10 ml of inoculum or in sterile MBR broth as a control for 30 min. Two inoculated seeds were planted in 600 g of soil/pot. Five ml of each bacterial suspension ( $\sim 1 \times 10^9$  CFU/ml/pot) were re-inoculated in the rhizosphere per treatment after 20 and 40 days of establishment. Soil chemical and physical properties are listed below: pH (7.15), organic matter (3.02%), effective cationic interchange coefficient (15.23 cmol/kg), P (215.93 ppm), S (15.08 ppm), Ca (13.37 cmol/kg), Mg (1.8 cmol/kg), K (0.63 cmol/kg), Na (0.05 cmol/kg), Fe (38.5 ppm), Mn (2.2 ppm), Cu (1.7 ppm). Plants grown at  $25\text{--}35^{\circ}\text{C}$ , 16:8 day/night regime with daily water application. After 90 days, flowering stage, root and shoot length were measured. Samples were dried at  $50^{\circ}\text{C}$  for 72 days and subsequently, dry weights were taken (boll, shoot and root). In addition, nitrogen (N) content was estimated over the total plant tissue in the LISSALAB Laboratory at Corpoica.

### Data analysis


Statistical analysis was performed using ANOVA ( $\alpha=0.05$ ) and the Tukey's HSD test. Associations among variables were examined by a simple correlation analysis. These analyses and graphics were performed using the GraphPad prism 7 software (Graphpad, San Diego, CA).

## Results and discussion

### Inoculation with bacteria improves plant growth regardless of chemical treatment

Rhizosphere is a highly dynamic environment in which complex interactions take place between plants and microorganisms. The understanding of those biological interactions represents a potential opportunity for biotechnological developments and could increase crop yield while reducing the use of agrochemicals<sup>28</sup>. In this study, the biofertilizer potential of *Azotobacter* strains with N fertilizer doses were evaluated in a glasshouse experiment to determine whether inoculation could replace N fertilization rates in cotton growth.

The results showed inoculation with bacteria caused a significantly increase ( $p < 0.05$ ) in plant growth compared to controls (Fig. 1). For example, when concentration of the chemical fertilizer was 50% of the recommended dosage, we observed inoculation with AC1, AC10 and their co-inoculation significantly ( $p < 0.05$ ) increased shoot length to



**Figure 1** Influence of *Azotobacter chroococcum* strains with different urea doses on growth parameters of cotton: shoot length (A), root length (B), shoot dry weight (C), root dry weight (D), boll dry weight (E), and N content (F). Media and standard errors are the result of six replicates per treatment and three independent experiments. Different letters indicate significant differences based on the HSD Tukey test.

9, 7 and 38% and root length to 6, 16 and 32%, respectively. In dry weight, root, shoot, and boll were significantly ( $p < 0.05$ ) enhanced to ~13, 20 and 25%, respectively. We also studied the effect of inoculation on N content. In general, microbial application led to a slight increase in N content. Therefore, significant improvements in N content by bacteria were only exhibited when mixed inoculation was applied, which indicates that bacterial inoculation has a greater influence on plant biomass respect to N content. The present findings suggest plant–bacteria interaction might be able to improve nutrient use efficiency of the fertilizer and/or bacterial inoculation might directly benefit plant growth<sup>1</sup>. It is important to note that the effect of AC1 was more pronounced on shoots than on roots, while AC10 had a greater effect on roots. A positive correlation ( $R = 0.65$ ) was observed between root length and N content on whole-plant tissue solely when AC10 was present. Kumar et al.<sup>18</sup>, who found that *A. chroococcum* had the ability to settle in cotton roots, support our findings by its chemotactic response

to root exudates. Furthermore, Adesemoye et al.<sup>3</sup>, proposed a combination of the activities of the plant and the inoculants as model for PGPR/PGPB-enhanced nutrient uptake in plants. We concluded that the increase in root length by PGPR/PGPB might contribute to an increased rate of N uptake in the plant.

### Co-inoculation enhances plant growth with respect to single inoculations

In this experiment, we observed that mixed inoculation with AC1 and AC10 increased plant growth with respect to single inoculations. Indeed, we evidenced an increase ranging between 8 and 53% in all parameters of plant growth, suggesting that microbial co-inoculation has a higher potential to promote plant growth (Fig. 1). The observed improvement in plant growth could be explained by a positive and additive contribution of each microorganism compared to single

Table 1 Multiple plant growth promotion traits *in vitro*

Strain	Nitrogenase activity		Solubilization				Phytohormulation			Production		
	ARA (nmol ethylene/ml h)		P (mg PO <sub>4</sub> <sup>-3</sup> /ml)	pH	Ca-P	Fe-P	pH	Musc	K	ACCd	IAA (mg indole/mg protein)	Siderop
AC1	68.01 ± 0.52	1408 ± 95.57	4.70 ± 0.07	76.33 ± 12.14	4.41 ± 0.06	-	-	-	-	0.16 ± 0.01	-	-
AC10	45.32 ± 1.94	1033 ± 47.75	4.79 ± 0.09	75.50 ± 15.71	4.34 ± 0.05	-	-	-	-	0.18 ± 0.01	-	-

Data are presented as mean ± standard error. Media and standard errors are the result of three replicates per biochemical analysis and results are representative of two independent experiments. (+/-) indicates presence/absence of trait. ARA: acetylene reduction assay; Ca-P: tricalcium phosphate; Fe-P: iron phosphate; Musc: muscovite; Biot: biotite; IAA: indole-3-acetic acid; ACCd: aminocyclopropane-1-carboxylate deaminase; Siderop: siderophore.

inoculations. Noteworthy, other investigators have observed similar results when using bacterial co-inoculations to promote plant growth<sup>2,9,20</sup>. Rojas-Tapias et al.<sup>30</sup>, in contrast to our observations, reported that co-inoculation did not lead to any additional effect compared to single inoculations of PGPR/PGPB when studying maize growth. This contrasting observation, however, might be associated to the evolutionary relationship among different plant and bacterial species, and soil chemical-physical properties<sup>26</sup>.

### Co-inoculation with *A. chroococcum* AC1 and AC10 strains allow to replace nitrogen fertilization

Notably, we found co-inoculation plus 50% urea caused a significantly effect on cotton growth in comparison with 75 and 50% of chemical fertilization, respectively. Furthermore, we evidenced 100% fertilization and 50% fertilization + co-inoculation produced similar results without statistical differences. In the presence of co-inoculation plus half urea doses, promotion of shoot length, shoot dry weight and boll dry weight were 10, 9 and 5% with respect to the complete fertilization program. In contrast, root length and N content were higher under full urea rates for 0.6 and 4%, respectively. This observation strongly suggests co-inoculation partially replaced the normal doses of nitrogen fertilization, which means that a reduction in 1.17 g/pot for urea or 37.5 kg per ha of urea is possible due to biological fertilization. Maheshwari et al.<sup>21</sup> reported similar findings, with *Sesamun indicum* showing a fertilizer reduction of 50% using *A. chroococcum* TRA2. In cotton, Pereg and McMillan<sup>24</sup>, described that the bacterial genera most commonly used to increase the efficiency of chemical fertilizers are *Azospirillum sp.*, *Methylobacterium sp.*, *Bacillus sp.*, *Pseudomonas sp.* and combination of *A. chroococcum* with arbuscular mycorrhizal fungi with comparable results on growth, yield and quality of fiber with respect to the complete fertilization program.

### AC1 and AC10 possess PGP traits

Bacterial inoculation led to increased plant growth. In order to understand the potential mechanisms involved in plant growth promotion, we characterized some of the PGP features of AC1 and AC10 using *in vitro* experiments. The results revealed that each strain exhibits more than one PGP attribute, which indirectly might explain the observed results in the glasshouse experiments. We observed that AC1 and AC10 were able to fix nitrogen, as expected in this bacterial genus. The observed values of acetylene reduction (AR) ranged between 1033 and 1408 nmol ethylene/ml h (Table 1). Interestingly, we found an inverse relationship among AR results and N content in total plant tissue in both microorganisms.

With respect to nutrient solubilization, we observed that strains were able to solubilize P from tricalcium phosphate at pH 7.5 and iron phosphate at pH 5.0. Negative correlations ( $R=0.88$ ) between pH and P solubilization were obtained in both P sources. This biological capability of strains was assessed under similar soil pH conditions and P insoluble sources. This result showed that AC1 and AC10 strains were more efficient in solubilizing Ca-P than Fe-P under our

**Table 2** Hydrolytic enzyme production

Strain	Hydrolytic enzyme synthesis					
	Protease	Cellulase	Pectinase	$\alpha$ -Amylase	Chitinase	Urease
AC1	+	–	–	–	–	+
AC10	+	–	+	–	–	+

Results are representative of two independent experiments. (+/–) indicates presence/absence of trait.

study conditions. Phosphate solubilization is associated to soil phosphorus availability increase, possibly conducted by the bacterial secretion of organic acids, acid phosphates and/or chelating agents into the growing medium<sup>23</sup>. However, negative results, were observed in K solubilization from muscovite and biotite.

Our results also demonstrated that none of the strains was able to produce iron siderophore complexes in CAS agar and did not possess ACC deaminase activity. To our knowledge, no evidence of the ability of *Azotobacter* to express ACCd activity has been not previously reported. Based on our observations, we associated root development with IAA production by the strains. Our cultures produced and excreted IAA to the medium in the presence of L-Tryptophan. Plant roots produce organic compounds including L-Tryptophan, which can be employed by PGPR/PGPB for IAA biosynthesis under natural conditions. IAA is the most physiologically active auxin in plants involved in cell enlargement and division, tissue differentiation, and responses to light and gravity<sup>32</sup>. Ultimately, we found production of hydrolytic enzymes such as proteases, urease, and pectinase in our strains (Table 2). These bacterial features not only confer ability to inhibit phytopathogens or to compete against soil microbial populations since their cell walls will be degraded by extracellular enzymes and their deleterious effects suppressed, but also contribute in the process of nutrient mineralization<sup>25</sup>.

Bacterial plant growth promotion is a well-established and complex phenomenon, which is often achieved by activities of more than one PGP characteristic exhibited by plant-associated bacteria<sup>27</sup>. In our study, both strains exhibited more than four PGP traits. Although several studies have focused on biological nitrogen fixation of *Azotobacter* as the primary influence on plant growth, these findings lead us to consider the possibility that improvement of cotton growth could be more related to other PGP activities of our strains than N uptake due to the ability to fix nitrogen.

We conclude that co-inoculation of *A. chroococcum* strains allow to reduce nitrogen fertilization doses up to 50% on cotton growth in its flowering stage under glasshouse conditions. Furthermore, biochemical analyses showed that both strains possess multiple PGP features, which indirectly might explain the observed results. Although our work does not establish whether bacterial mechanisms were responsible by the positive effect of co-inoculation, this study provides a starting point for future research in which molecular tools in different metabolic pathways will allow to discern the mechanisms associated with the results reported here. Therefore, field tests should be carried out to obtain information regarding the response of cotton plants in terms of yield and cotton fiber quality, when inoculated with AC1

and AC10 strains. We consider that a reduction in the amount of agrochemicals used in cotton crop is paramount to reduce its impact on environmental health and contribute to the mitigation of climate change.

## Ethical disclosures

**Protection of human and animal subjects.** The authors declare that no experiments were performed on humans or animals for this study.

**Confidentiality of data.** The authors declare that no patient data appear in this article.

**Right to privacy and informed consent.** The authors declare that no patient data appear in this article.

## Funding

This work was financed by Ministerio de Agricultura y Desarrollo Rural of Colombia.

## Conflict of interest

The authors declare that they have no competing interest in relation to work presented in this manuscript.

## Acknowledgements

We thank to Mauricio Barón for the support in the glasshouse experiment, Daniel Bravo for help with determining K solubilization, Jorge Arguelles for statistical analysis, and Diego Rincón and Juan Pablo Hernández for critical review on the manuscript.

## References

- Adesemoye AO, Torbert HA, Kloepper KW. Enhanced plant nutrient use efficiency with PGPR and AMF in an integrated nutrient management system. *Can J Microbiol.* 2008;54:876–86.
- Adesemoye AO, Kloepper JW. Plant–microbes interactions in enhanced fertilizer-use efficiency. *Appl Microbiol Biotechnol.* 2009;85:1–12.
- Adesemoye AO, Torbert HA, Kloepper JW. Plant growth-promoting rhizobacteria allow reduced application rates of chemical fertilizers. *Microb Ecol.* 2009;58:921–9.
- Bhattacharyya PN, Jha DK. Plant growth-promoting rhizobacteria (PGPR): emergence in agriculture. *World J Microbiol Biotechnol.* 2012;28:1327–50.

5. Bonilla R, Novo R, Roncallo B, Camelo M, Criollo P, Estrada GA, Garrido MF, López M, Moreno-Galván A, Obando M, Parra L, Rivera D, Rojas-Tapias D, Van Strahlen M, Zambrano C. *Azotobacter* sp. applied research aimed to growth promotion of crops with economics interest in Colombia. In: García de Salomone IE, Vásquez S, Penna C, Cassán F, editors. *Rizósfera, Biodiversidad y Agricultura Sustentable*. 1st ed. Argentina: Lit Verlag; 2013. p. 151–73.
6. Bradford MM. A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Anal Biochem*. 1976;72:248–54.
7. Cappuccino JG, Sherman N. *Microbiology: a laboratory manual. The Benjamin/Cummings Publishing Co.*; 2014. ISBN13: 9780321840226.
8. Choudhary DK, Kasotia A, Jain S, Vaishnav A, Kumari S, Sharma KP, Varma A. Bacterial-mediated tolerance and resistance to plants under abiotic and biotic stresses. *J Plant Growth Regul*. 2016;35:276–300.
9. Compant S, Clément C, Sessitsch A. Plant growth-promoting bacteria in the rhizo- and endosphere of plants: their role, colonization, mechanisms involved and prospects for utilization. *Sol Biol Biochem*. 2010;42:669–78.
10. De Bolle S, Gebremikael MT, Maervoet V, De Neve S. Performance of phosphate solubilizing bacteria in soil under high phosphorous conditions. *Biol Fertil Soils*. 2013;49:705–14.
11. Eckert B, Weber OB, Kirchhof G, Halbritter A, Stoffels M, Hartmann H. *Azospirillum doebereineriae* sp. nov. a nitrogen-fixing bacterium associated with the C4-grass *Miscanthus*. *Int J Syst Evol Microbiol*. 2001;51:17–26.
12. Glick BR. Plant growth-promoting bacteria: mechanisms and applications. *Scientifica*. 2012. <http://dx.doi.org/10.6064/2012/963401>, 15 pp. Article ID 963401.
13. Glick BR. *Beneficial plant–bacterial interactions*. Springer International Publishing; 2015. p. 5. <http://dx.doi.org/10.1007/978-3-319-13921-0>.
14. Glickmann E, Dessaux Y. A critical examination of the specificity of the salkowski reagent for indolic compounds produced by phytopathogenic bacteria. *Appl Environ Microbiol*. 1995;61:793–6.
15. Gurikar C, Naik MK, Sreenivasa MY. *Azotobacter*: PGPR activities with special reference to effect of pesticides and biodegradation. In: Singh MK, Singh HB, Prabha R, editors. *Microbial inoculants in sustainable agricultural productivity*. 1st ed. India: Springer; 2016. p. 229–44.
16. Honna M, Shimomura T. Metabolism of 1-aminocyclopropane-1-carboxylic acid. *Agric Biol Chem*. 1978;42:1825–31.
17. Hu XF, Chen J, Guo JF. Two phosphate- and potassium-solubilizing bacteria isolated from Tianmu Mountain, Zhejiang, China. *World J Microbiol Biotechnol*. 2006;22:983–90.
18. Kumar R, Bhatia R, Kukreja K, Behl RK, Dudeja SS, Narula N. Establishment of *Azotobacter* on plant roots: chemotactic response, development and analysis of root exudates of cotton (*Gossypium hirsutum* L.) and wheat (*Triticum aestivum* L.). *J Basic Microbiol*. 2007;47:436–9.
19. Kraiser T, Gras DE, Gutierrez AG, González B, Gutiérrez RA. A holistic view of nitrogen acquisition in plants. *J Exp Bot*. 2011;62:1455–66.
20. Lavakush YJ, Verma JP, Jaiswal DK, Kumar A. Evaluation of PGPR and different concentration of phosphorus level on plant growth, yield and nutrient content of rice (*Oryza sativa*). *Ecol Eng*. 2014;62:123–8.
21. Maheshwari DK, Dubey RC, Aeron A, Kumar B, Kumar S, Tewari S, Arora NK. Integrated approach for disease management and growth enhancement of *Sesamun indicum* L. utilizing *Azotobacter chroococcum* TRA2 and chemical fertilizer. *World J Microbiol Biotechnol*. 2012;28:3015–24.
22. Moreno-Galván A, Rojas-Tapias D, Bonilla R. Sequential statistical design application in identification of *Azotobacter chroococcum* AC1 nutritional sources. *Corpoica Cienc Tecnol Agropecu*. 2011;11:151–8.
23. Panhwar QA, Naher UA, Jusop S, Othman R, Latif MA, Ismail MR. Biochemical and molecular characterization of potential phosphate-solubilizing bacteria in acid sulfate soils and their beneficial effects on rice growth. *PLoS ONE*. 2014;9:e97241.
24. Pereg L, McMillan M. Scoping the potential uses of beneficial microorganisms for increasing productivity in cotton cropping systems. *Soil Biol Biochem*. 2015;80:349–58.
25. Pereira SI, Castro PM. Phosphate-solubilizing rhizobacteria enhance *Zea mays* growth in agricultural P-deficient soils. *Ecol Eng*. 2014;73:526–35.
26. Prosser JI, Bohannan BJ, Curtis TP, Ellis RJ, Firestone MK, Freckleton RP, Green LE, Killham K, Lennon JK, Osborn AM, Solan M, Van der Gast CJ, Young PW. The role of ecological theory in microbial ecology. *Nat Rev Microbiol*. 2007;5:384–92.
27. Rana A, Saharan B, Joshi M, Prasanna R, Kumar K, Nain L. Identification of multitrait PGPR isolates and evaluating their potential as inoculants for wheat. *Ann Microbiol*. 2011;61:893–900.
28. Roca A, Pizarro-Tobías P, Udaondo Z, Fernández M, Matilla MA, Molina-Henares MA, Molina L, Segura A, Duque E, Ramos JL. Analysis of the plant growth-promoting properties encoded by the genome of the rhizobacterium *Pseudomonas putida* BRID-1. *Environ Microbiol*. 2013;15:780–94.
29. Rojas-Tapias D, Moreno-Galván A, Pardo-Díaz S, Obando M, Rivera D, Bonilla R. Effect of inoculation with plant growth-promoting bacteria (PGPB) on amelioration of saline stress in maize (*Zea mays*). *Appl Soil Ecol*. 2012;61:264–72.
30. Rojas-Tapias D, Bonilla R, Dussán J. Effect of inoculation and co-inoculation of *Acinetobacter* sp. RG30 and *Pseudomonas putida* GN04 on growth, fitness, and copper accumulation of maize (*Zea mays*). *Water Air Soil Pollut*. 2014;225:1–13.
31. Schwyn B, Neilands JB. Universal chemical assay for the detection and determination of siderophores. *Anal Biochem*. 1987;160:47–56.
32. Sukumar P, Legue V, Vayssieres A, Martin F, Tuskan GA, Kalluri UC. Involvement of auxin pathways in modulating root architecture during beneficial plant–microorganism interactions. *Plant Cell Environ*. 2013;36:909–19.
33. Zhang X, Davidson EA, Mauzerall DL, Searchinger TD, Dumas P, Shen Y. Managing nitrogen for sustainable development. *Nature*. 2015;528:51–9.