



MICROBIOLOGICAL IMAGE

Nematicidal effect of an Argentine strain of *Photorhabdus laumondi laumondi* (Enterobacteriaceae) on the free-living nematode *Panagrellus redivivus* (Rhabditidae: Panagrolaimidae)

Efecto nematocida de una cepa argentina de *Photorhabdus laumondi laumondi* (Enterobacteriaceae) sobre el nematodo de vida libre *Panagrellus redivivus* (Rhabditidae: Panagrolaimidae)

Augusto Salas^a, María Fernanda Achinelly^b, Diego Herman Sauka^{a,*}

^a Instituto Nacional de Tecnología Agropecuaria (INTA), Instituto de Microbiología y Zoología Agrícola, Argentina

^b Centro de Estudios Parasitológicos y de Vectores (CEPAVE), Facultad de Ciencias Naturales y Museo, Universidad Nacional de La Plata, Argentina

Received 23 December 2019; accepted 6 April 2020

Available online 31 May 2020

Bacteria with nematicidal effects have attracted interest as biocontrol agents of pests and diseases due to their relatively low environmental impact compared to chemical management strategies¹. The damage they cause to target nematodes is diverse and imperfectly understood⁴.

Photorhabdus spp. (Enterobacteriaceae) lives symbiotically within the nematode *Heterorhabditis* spp. (Rhabditida: Heterorhabditidae). It is known that secondary metabolites of this gram-negative bacterium exhibit insecticidal activity, mainly in Lepidopteran and Coleopteran insects². Studies have also shown that cell-free culture filtrates of *Photorhabdus* spp. have nematicidal activity on free-living nematodes

and pest phytonematodes; however, the causes of death in these targets are still being studied³.

This work provides information on the effect caused by the intake of *Photorhabdus laumondi laumondi* (strain LP1900, isolated from *Heterorhabditis bacteriophora* strain SUP2), in the digestive tract of the bacteriophagous nematode *Panagrellus redivivus* (Rhabditidae: Panagrolaimidae).

LP1900 strain was cultured on nutrient agar (Difco) plates for 7 days at 29 °C (5 repetitions); then 2×10^3 nematodes were inoculated on each plate, and freely dispersed through the culture. Untreated controls consisted of agar nutrient plates with the 2×10^3 nematode inoculum alone. Alterations in the posterior esophagus (behind the basal bulb), and in the anterior, middle and posterior intestinal regions of treated nematodes were recorded 24 h after treatment under a light microscope (10×) (Eclipse E200; Nikon), and compared with untreated nematodes (Fig. 1).

* Corresponding author.

E-mail address: sauka.diego@inta.gob.ar (D.H. Sauka).




Figure 1 Nematicidal effect of *P. laumondi laumondi* LP1900 on the digestive tract of *P. redivivus*. (A) Both the intestine and the cuticle were intact in untreated nematodes. (B) A rupture in the cuticle near the basal bulb region can be observed on nematodes exposed to a 7-day-old culture of LP1900, spilling the internal contents of the organism. The musculature, from the stoma to the beginning of the basal bulb, showed signs of disintegration. (C, D) A similar effect was observed in the middle and posterior intestinal regions of other treated nematodes.

Acknowledgments

AS worked with fellowship from Consejo Nacional de Investigaciones Científicas y Técnicas (CONICET). DS and MFA are members of the Research Career of CONICET.

We would like to thank Mariano Maestro for the English revision of the text.

References

1. Bojanich M, Basualdo J, Giusiano G. In vitro effect of *Chrysosporium indicum* and *Chrysosporium keratinophyllum* on *Toxocara canis* eggs. *Rev Argent Microbiol.* 2018;50:249–54.
2. Eliceche D, Tesis Estudio de las características bio-ecológicas de un aislamiento autóctono del entomonemátodo *Heterorhabditis bacteriophora* (Nematoda: Heterorhabditidae) y su potencialidad como agente bioregulador de insectos. Buenos Aires, Argentina: Facultad de Ciencias Naturales y Museo, UNLP; 2019.
3. Hu K, Li J, Webster J. Nematicidal metabolites produced by *Photorhabdus luminescens* (Enterobacteriaceae), bacterial symbiont of entomopathogenic nematodes. *Nematology.* 1999;1:457–69.
4. Maestro M, Sauka D. Progression of the predatory activity of an Argentine strain of *Arthrobotrys oligospora* (Helotiales: Orbiliaceae) on *Panagrellus redivivus* (Rhabditidae: Panagrolaimidae). *Rev Argent Microbiol.* 2019, <http://dx.doi.org/10.1016/j.ram.2019.03.008>.