



MICROBIOLOGICAL IMAGE

Endophytic microorganisms *Agrobacterium tumefaciens* 6N2 and *Meyerozyma guilliermondii* 6N serve as models for the study of microbial interactions in colony biofilms



Los microorganismos endofíticos *Agrobacterium tumefaciens* 6N2 y *Meyerozyma guilliermondii* 6N sirven como modelos para el estudio de las interacciones microbianas en *biofilms* en colonia

Elisa Violeta Bertini^a, Ana Carolina del V. Leguina^a,
Lucía I. Castellanos de Figueroa^{a,b}, Carlos G. Nieto-Peñalver^{a,b,*}

^a PROIMI, CONICET (Planta Piloto de Procesos Industriales Microbiológicos), Tucumán, Argentina

^b Instituto de Microbiología, Facultad de Bioquímica, Química y Farmacia, Universidad Nacional de Tucumán, Tucumán, Argentina

Received 18 June 2018; accepted 7 September 2018

Available online 14 January 2019

We analyzed the morphology of colonies of endophytic bacteria and yeasts by fluorescence. *Agrobacterium tumefaciens* 6N2 (formerly 197MX) is a non-pathogenic isolate from sugarcane with features of relevance for microbial interactions¹. *Meyerozyma guilliermondii* 6N was obtained from the same host³.

A. tumefaciens 6N2 was tagged with *gfp.AAV-a*, allowing the detection of cells actively expressing the fluorescence². GFP-tagged 6N2 and *M. guilliermondii* 6N were cultured at 30 °C in nutrient broth until late exponential phase¹. Cell densities were adjusted to ~0.1 OD_{600nm}, and spot inoculated on nutrient agar (NA) and YPD agar plates⁵, as pure or mixed cultures (proportion 1:1). Plates were incubated at 30 °C for 72 h; colonies were imbibed with 10 μl of 25 μM

Calcofluor White M2R (CW) that binds chitin and cellulose, and observed with a 4× objective lens under a Zeiss SterEO Lumar.V12 Stereomicroscope with GFP and DAPI filters.

While 6N2 colonies exhibited a wrinkled surface on YPD agar (Fig. 1A and C), they were smooth and convex on NA (Fig. 2A and C). Under interaction conditions 6N2 showed singular patterns: 6N2 was located at the edges and the center on YPD agar (Fig. 1D), and was evenly distributed on NA (Fig. 2D). In both media 6N seemed to be located on top of the bacterial growth (Fig. 1B and B). 6N development was granular with lobate borders on YPD agar (Fig. 1B and F), and was smooth and dotted on NA (Fig. 2B and F). Noteworthy, 6N2 was not stained with CW (Fig. 1E and E). Beyond the effect of the interactions on the host plant, the images show

* Corresponding author.

E-mail address: cgneto@proimi.org.ar (C.G. Nieto-Peñalver).




Figure 1 Colony biofilms of *A. tumefaciens* 6N2 and *M. guilliermondii* 6N growing on YPD agar. A, C and E: pure colony of 6N2 observed with visible illumination, GFP filter and DAPI filter, respectively. B, D and F: colony of mixed cultures of 6N2 and 6N observed with visible illumination, GFP filter and DAPI filter, respectively.




Figure 2 Colony biofilms of *A. tumefaciens* 6N2 and *M. guilliermondii* 6N growing on NA. A, C and E: pure colony of 6N2 observed with visible illumination, GFP filter and DAPI filter, respectively. B, D and F: colony of mixed cultures of 6N2 and 6N observed with visible illumination, GFP filter and DAPI filter, respectively.

the potential of the strains and the techniques for the study of interactions in a colony biofilm, even if the true location of each strain requires other techniques (e.g., confocal microscopy)⁴.

Funding

This work was supported by the Consejo Nacional de Investigaciones Científicas y Técnicas (CONICET, PIP 946, PU-E 22920160100012CO), Agencia Nacional de Promoción Científica y Tecnológica (PICT 2016 No. 0532; PICT 2016 No. 2013), and Secretaría de Ciencia, Arte e Innovación Tecnológica from the Universidad Nacional de Tucumán (PIUNT D609).

Conflict of interest

The authors declare that they have no conflicts of interest.

Acknowledgements

We thank Jean-Michel Camadro (Institut Jacques Monod, UMR7592, CNRS Université Paris Diderot, Paris, France) for kindly providing access to the stereomicroscope facility.

References

1. Bertini EV, Leguina AC, del V, Castellanos de Figueroa LI, Nieto Peñalver CG. Caracterización de bacterias endofíticas de caña de azúcar productoras de *N*-acil homoserina lactonas. *Archivos de Bioquímica, Química y Farmacia*. 2016;25:5–19.
2. Lamberts L, Sternberg C, Molin S. Mini-Tn7 transposons for site-specific tagging of bacteria with fluorescent proteins. *Environ Microbiol*. 2004;6:726–32.
3. Leguina AC, del V, Nieto C, Pajot HM, Bertini EV, Mac Cormack W, Castellanos de Figueroa LI, Nieto-Peñalver CG. Inactivation of bacterial quorum sensing signals *N*-acyl homoserine lactones is widespread in yeasts. *Fungal Biol*. 2018;122:52–62.
4. Merritt JH, Kadouri DE, O'Toole GA. Growing and analyzing static biofilms. In: *Curr Protoc Microbiol*; 2005. pp. 1B.1.1–1B.1.18.
5. Yeast Extract–Peptone–Dextrose (YPD) Medium (Liquid or Solid). In: *Cold Spring Harb Protoc*; 2017, <http://dx.doi.org/10.1101/pdb.rec090563>.