



ORIGINAL ARTICLE

Isolation of culturable mycota from Argentinean soils exposed or not-exposed to pesticides and determination of glyphosate tolerance of fungal species in media supplied with the herbicide



Melisa Egle Aluffi, Cecilia Soledad Carranza, Nicolás Benito, Karen Magnoli, Carina Elizabeth Magnoli, Carla Lorena Barberis*

Instituto de Investigación en Micología y Micotoxicología (IMICO-CONICET), Departamento de Microbiología e Inmunología, Facultad de Ciencias Exactas, Físico, Químicas y Naturales, Universidad Nacional de Río Cuarto, Ruta Nacional N° 36 Km 601, 5800 Río Cuarto, Córdoba, Argentina

Received 31 July 2019; accepted 11 November 2019

Available online 9 January 2020

KEYWORDS

Culturable mycota;
Soils;
Glyphosate;
Tolerance;
Growth parameters

Abstract The current agricultural system has led to the development of glyphosate (GP)-resistant weeds, causing an increase in GP doses and applications. Native mycota of pesticide-contaminated sites are the major source of pesticide-degrading microorganisms. The aims of the present study were to isolate the GP-tolerant culturable mycota in two soils with different pesticide exposure from Córdoba, Argentina, and to evaluate the growth parameters in native fungal isolates in the presence of GP and the effective dose that caused 50% growth reduction. The results showed that the genera *Fusarium*, *Aspergillus*, *Mucor*, *Penicillium* and *Sterilia* were the prevalent fungi isolated from soils both exposed and not-exposed to pesticides. The highest value (>100 mM) of effective concentration of herbicide that caused 50% growth inhibition (EC50), was found for *Trichoderma* isolates. *Sterilia* spp. had EC50 values of 100 mM, while *Aspergillus* spp. and *Mucor* had EC50 values between 50 and 100 mM. The growth rate evaluation varied according to the isolates and GP concentrations. The data showed that all *Aspergillus* spp., *Trichoderma* spp., *Mucor* and three *Sterilia* spp. had the best growth performance in media supplied with GP after a variable acclimation period. This study provides valuable data for further studies that would allow to know the metabolic capacity of these fungal species that can be potential candidates for GP removal from contaminated environments.

© 2019 Asociación Argentina de Microbiología. Published by Elsevier España, S.L.U. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

* Corresponding author.

E-mail address: cbarberis@exa.unrc.edu.ar (C.L. Barberis).

PALABRAS CLAVE

Microbiota cultivable;
Suelos;
Glifosato;
Tolerancia;
Parámetros de
crecimiento

Aislamiento de la microbiota cultivable de suelos argentinos expuestos y no expuestos a plaguicidas y determinación de la tolerancia a glifosato de especies fúngicas en medios suplementados con el herbicida

Resumen El sistema agrícola actual condujo al desarrollo de malezas resistentes a glifosato (GP), lo que deriva en el aumento de las dosis y el número de aplicaciones requeridas. La microbiota nativa de sitios contaminados con plaguicidas constituye la fuente principal de microorganismos capaces de degradar este herbicida. Los objetivos del presente estudio fueron aislar la microbiota cultivable tolerante a GP a partir de dos suelos agrícolas con diferente grado de exposición a plaguicidas, ambos de la provincia de Córdoba, Argentina, y evaluar en los hongos nativos aislados los parámetros de crecimiento en presencia de GP y la dosis efectiva que produce un 50% de reducción del crecimiento (EC50). Los géneros *Fusarium*, *Aspergillus*, *Mucor*, *Penicillium* y *Sterilia* fueron los de mayor prevalencia, tanto en suelos expuestos como no expuestos a plaguicidas. La mayor EC50 hallada (> 100 mM) correspondió a dos aislamientos de *Trichoderma*. En dos aislamientos de *Sterilia*, los valores de EC50 fueron de 100 mM, mientras que en los dos de *Aspergillus* y en uno de *Mucor* fueron de entre 50 y 100 mM. La velocidad de crecimiento varió según la concentración de GP empleada y el aislamiento considerado. Todos los aislamientos de *Aspergillus*, *Trichoderma* y *Mucor*, y tres aislamientos de *Sterilia* mostraron el mejor rendimiento en cuanto al crecimiento en los medios con GP después de un período de aclimatación variable. Este trabajo proporciona datos preliminares valiosos referidos a la capacidad metabólica de estos aislados y su potencial para remover GP en ambientes contaminados.

© 2019 Asociación Argentina de Microbiología. Publicado por Elsevier España, S.L.U. Este es un artículo Open Access bajo la licencia CC BY-NC-ND (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Introduction

Herbicides based on glyphosate (HBGs) can be applied several times throughout the year, even when the crops are growing or between growing seasons in order to hasten the drying of weeds³⁰. In Argentina this agricultural system has led to the development of GP-resistant weeds, causing an increase in both GP doses and applications. Therefore, it was necessary to apply different formulations of herbicides to achieve the control of all weeds. The Humid Pampa region of Argentina is responsible for 80% of soybean production with intensive use of agrochemicals, HBGs being the most applied herbicides³⁵.

Recent studies have revealed that N-phosphonomethylglycine (GP) and its main degradation metabolite aminomethylphosphonic acid (AMPA) may persist in soil for long periods⁸. The residues are found at all levels of the food chain, such as drinking water, plants, animals, and even in exposed humans³⁴. The presence of GP and AMPA has been reported in water and sediments of streams from rural and suburban basins of our country within the provinces of Buenos Aires, Santa Fe and Córdoba^{2,41,43,18}. The extensive use of this herbicide has shown to cause a selection in soil microbial populations, with increases in populations of specific microbial taxa with capacity to degrade the herbicide³¹.

The reports of International organizations such as the Food and Agricultural Organization of the United Nations have alerted about pesticide pollution²¹. In this context, several researchers worldwide have focused their attention on the development of strategies for minimizing the

pesticide contamination of natural resources and remediating contaminated natural environments³⁷. The bioaugmentation of contaminated sites with degrading microorganisms is one of the most promising bioremediation strategies. The soil is the main receiver of pesticides used in agriculture. After the application of HBG, a part of the herbicide is adsorbed or immobilized to the soil particles and the rest can leach into other ecosystems. The permanence of GP not linked to the soil depends on the photo, chemical, and microbial degradation rate. Bacteria and fungi are considered to be the most important microorganisms involved in biotic pesticide degradation. Mycelial growth and extracellular enzymes provide filamentous fungi an advantage over other microorganisms such as bacteria and yeasts. Furthermore, in acidic medium, soil fungal degradation prevails in the proximity of the surface where the oxygen and the organic matter is high⁶. In this context, fungal enzymes are important in biotechnology processes and are promising in organophosphorus herbicide degradation. These processes include mineralization, co-metabolism and interspecific coordination metabolism³⁷. Several reports have reported the ability of native soil microfungi such as *Aspergillus*, *Penicillium* and *Fusarium* in organophosphorus pesticide degradation⁶. In previous studies Carranza et al.^{15,12,13,16} showed that non-toxigenic *Aspergillus* section *Flavi* strains isolated from soil exposed to pesticides were able to tolerate *in vitro* high doses of GP (500 mM), and were successfully able to use the herbicide as both phosphorus or nitrogen source. The degradation percentages were higher than 50% at 15 days of incubation. These strains were tested on soil microcosms showing GP tolerance, permanence and

competitiveness in the presence of native mycota. Considering that the contaminated sites are the major source of degrading microorganisms and degradation of pesticides is not performed by a single microbial population^{6,37}, the aims of the present study were to isolate the GP-tolerant culturable mycota in two soils with different pesticide exposure, from Córdoba, Argentina, and to evaluate the growth parameters in native fungal isolates in the presence of GP and the effective dose that caused 50% of growth reduction.

Materials and methods

Sampling sites and sample processing

Four fields were selected for soil sampling in the south of the Province of Córdoba. Three fields were chosen with a long history of exposure to pesticides (approximately 10 years). From each field, three samplings (10 samples were taken in each sampling) along the year were performed between 2016 and 2017. The fields were located in: place 1 – Coronel Moldes (34°28'18" 63°32'18"), place 2 – Serrano (33°37'00" 64°30'23") and place 3 – Espinillo (33°00'47" 64°19'13"). Additionally, representative soil samples from fields without direct exposure to pesticides, no-tillage and with native forests located in Córdoba, Argentina (place 4: 31°56'16" 64°37'11") were analyzed. Twenty samples were collected from them in 2017.

Soil samples of 1 kg were collected from the surface layer (at a depth of 10 cm) of the soil. These samples were homogenized and air-dried for 1–2 days at 25–30 °C. Samples weighing 100 g were thoroughly mixed and passed through a testing sieve (2 mm mesh size) to separate the soil from the debris. Samples were stored at 4 °C and the isolation of fungi was performed within 2 days after the sampling.

Glyphosate solutions

A commercial formulation (Roundup Controlmax®, Buenos Aires, Argentina) was used in this study. A stock herbicide solution of 2 M of the active ingredient was prepared dissolving the appropriate quantity of commercial product in sterile distilled water (100 ml). The herbicide solution was sterilized (filters of 0.2 µm, Microclar, Buenos Aires, Argentina) and maintained at 4 °C until use.

Enrichment and isolation of fungi in media supplied with GP as carbon, nitrogen and phosphorous sources

Ten grams of each soil sample were added in Erlenmeyer flasks containing 50 ml of sterile modified Czapeck-Dox medium (CZD), with the following composition (l); 10 g glucose, 0.5 g MgSO₄·7H₂O, 0.5 g KCl, 2 g NaNO₃, 0.01 g FeSO₄, 1 g KH₂PO₄, 0.5 g yeast extract, gentamicin and chloramphenicol (100 mg/l each) to inhibit bacterial growth. CZD without glucose, NaNO₃ or KH₂PO₄; and supplied with GP at a final concentration of 10 mM (CZDC); 1.5 mM (CZDN) and 1.0 mM (CZDP), respectively were prepared to evaluate the use of GP as sole source of carbon, nitrogen and phosphorus³³. The CZD media were autoclaved at 121 °C for

15 min and the appropriate aliquot of GP was added to the sterilized culture media template at 45–50 °C to obtain the final concentrations mentioned above. These concentrations were selected based on the percentages that the carbon, nitrogen or phosphorus sources should be present in the fungal media. Each condition (CZDG, CZDN and CZDP) was evaluated by duplicate. Then, the flasks were incubated at 28 °C on a shaking platform at 120 rpm for 7 days. After the incubation period, serial dilutions from 10⁻¹ to 10⁻⁶ were performed on 0.1% peptone water solution and then aliquots of 0.1 ml were inoculated by the surface-spray method by duplicate on Petri plates containing CZDC, CZDN and CZDP media with 1.5% agar-agar. The plates were incubated in the dark at 28 °C for 7 days. After the incubation period, the plates that contained 10–100 colonies were used for total and genera counting.

Morphological identification at the genus level of the different colonies was conducted by sub-culturing each colony on Malt Extract Agar (MEA). The identification was performed through macroscopic and microscopic criteria following taxonomic keys^{42,46,49}. The results were expressed as colony-forming units (CFU) per gram of soil of the total mycota and each genus in each medium supplied with GP. The strains were incorporated into the culture collection of the Department of Microbiology and Immunology, National University of Río Cuarto, Córdoba, Argentina, and are maintained in 15% glycerol (Sigma Aldrich, St. Louis, MO, USA).

In vitro screening of GP tolerance of fungal isolates at optimal water availability and temperature conditions

All the strains that were isolated from the soils either exposed or not exposed to pesticides were subjected to three successive cultures in MEA supplied with 10 mM of GP for screening their capacity to grow in the presence of the herbicide. Of these, eighteen fungal isolates belonging to *Aspergillus*, *Trichoderma*, *Mucor*, *Fusarium* and *Penicillium* genera were selected to evaluate GP tolerance. All strains were isolated from exposed soils, except for *Trichoderma* spp. (strain 311). Lag phase prior to growth, growth rate and the effective dose that caused 50% of growth rate reduction (EC50) in the presence of GP were evaluated in order to estimate GP tolerance.

The CZD medium supplied with the solution of GP at final concentrations of 5, 20, 50 and 100 mM (equivalent to 0.85, 3.4, 8.4, and 16.9 mg/ml, respectively) was used for this assay. The lowest GP concentrations tested are the field application rates recommended and the highest ones represent the concentrations of pesticide reported in sites with spill³⁸. The water activity (*a_w*) of the medium was conditioned at 0.98 with glycerol to simulate the optimal conditions of growth. The *a_w* of the medium was checked with AquaLab Series 3 (Decagon Devices, Inc., Pullman, WA, USA). In addition, the respective controls without GP were prepared. The culture plates for each treatment were needle-inoculated centrally with a spore suspension (10⁶ spores/ml). The inoculum was prepared from 5- or 7-day-old cultures on MEA medium. Each treatment was done in triplicate and incubated at 25 °C for 15 days or until the

fungal colony reached the edge of the plate. All the experiment was repeated twice.

Growth rate was calculated taking two measures of colony diameter at right angles to one another daily from each replicate plate. The radius of the colony was plotted against time, and a linear regression was applied in order to obtain the growth rate as the slope of the line to the X-axis. The lag phase (h) before exponential growth was also determined⁷. The percentage of growth inhibition by GP was calculated for each isolate in each treatment in order to determine the EC50⁴⁰.

Statistical analysis

Data from the fungal count, growth rate and lag phases were subjected to analysis of variance. Means were compared using a linear mixed model and the Fisher's protected least significant difference (LSD) test to determine the significant differences between the means from total fungal counts and frequency of fungal genera. All data were transformed to $\log_{10}(x+1)$ to obtain homogeneity of variances. Means were also compared using the Fisher's protected LSD test to determine the influence of concentration of herbicide on the growth rate and lag phase prior to growth of the tested strains. The statistical analysis was performed using InfoStat Professional software version 2017²⁰.

Results

Enrichment and isolation of fungi in media supplied with GP as carbon, nitrogen and phosphorous sources

Figure 1 shows the mean values of total culturable mycota counts (\log_{10} CFU/g) from different sampling places in control media and supplied with GP. Total fungal counts varied according to sampling place. Counts ranged between 5.5 and 8.7 \log_{10} CFU/g. The highest values were observed in soil samples without pesticide exposure or no-tillage (place 4) ($p < 0.05$). In this place the highest counts were found in control media, followed by CDZN, CZDP and CDZC media. While in exposed places, the highest counts were found in media supplied with GP, and the counts varied according to the media supplied with GP analyzed.




Figure 1 Mean values of total fungal counts (\log_{10} CFU/g) from the different places sampled in control media (CZD) and supplied with GP as only source of nitrogen (CZDN), phosphorus (CZDP) and carbon (CZDC). ^{a,f}Mean values with different letters indicate significant differences in accordance with the Fisher's LSD test ($p < 0.05$).

With regard to the analysis of the culturable mycota, 13 genera of filamentous fungi were isolated. *Fusarium* spp., *Aspergillus* spp., *Mucor* spp., *Penicillium* spp., *Trichoderma* spp., *Paecilomyces* spp., *Cladosporium* spp., *Verticillium* spp., *Exophiala* spp., *Alternaria* spp., *Phoma* spp., *Phialophora* spp. and *Streptothrix* spp. were isolated in all soil samples. Some genera, such as *Alternaria* spp. and *Streptothrix* spp. were found only in non-exposed soil samples, with mean counts of 7 and 6 \log_{10} CFU/g, respectively. The genera *Trichoderma* spp., *Paecilomyces* spp. and *Verticillium* spp. were found only in the soils exposed to pesticides, with mean counts of 6.4, 5.4 and 6.3 \log_{10} CFU/g, respectively (data not shown).

Figure 2 shows the counts of prevalent genera from different places in control and GP-supplied media as sole source of nitrogen, phosphorous and carbon. The genera *Fusarium* spp., *Aspergillus* spp., *Mucor* spp., *Penicillium* spp. and *Sterilia* spp. were the prevalent fungi isolated from soils both exposed and not-exposed to pesticides. They were isolated in all the media tested, except *Mucor* spp., which were only isolated in control and CZDP medium without significant differences between both media. A different behavior was observed in the other genera. *Fusarium* spp. showed similar counts in control and CZDN and CZDP media. For *Aspergillus* spp. the highest count was found in the CZDP medium. While for *Penicillium* spp. this behavior was observed in CZDP and CZDN. For *Sterilia* spp. no significant differences in the count were found among the supplied GP media ($p < 0.05$).

In vitro screening of GP tolerance of fungal isolates at optimal water availability and temperature conditions

Table 1 shows the lag phase prior to growth and EC50 of 18 isolates in media with increasing GP concentrations at optimal water availability and temperature conditions. Lag phase varied depending on the isolate tested. In general, the presence of GP in the media produced a time of adaptation similar or higher than those in the control media. For most of the isolates, the highest GP concentration (100 mM) produced longer lag phases. Only one isolate (*Penicillium* spp., 139) did not develop in this condition (the lag phase was extended more than the




Figure 2 Mean values (\log_{10} CFU/g) of the prevalent genera from different places in control media and supplied with GP as only source of nitrogen (CZDN), phosphorus (CZDP) and carbon (CZDC). ^{a,e}Mean values with different letters indicate significant differences in accordance with the Fisher's LSD test ($p < 0.05$).

Table 1 Effect of different concentrations of GP on the lag phase (h) and EC50 of different isolates in CZD medium at 0.98 a_w and 25 °C.

Isolate	Genera	Lag phase (h) \pm SD Glyphosate (mM)					EC50
		0	5	20	50	100	
128	<i>Aspergillus</i> spp.	30.1 \pm 3.6 ^a	29.3 \pm 5.6 ^a	25.4 \pm 21.5 ^a	32.9 \pm 17.2 ^a	40.3 \pm 35.0 ^b	50–100
2	<i>Aspergillus</i> spp.	30.0 \pm 1.0 ^a	26.5 \pm 0.7 ^a	28.9 \pm 0.3 ^a	19.4 \pm 18.3 ^a	20.0 \pm 5.3 ^a	50–100
133	<i>Sterilia</i> spp.	49.0 \pm 1.7 ^{ab}	43.8 \pm 18.7 ^{ab}	56.2 \pm 6.1 ^{ab}	54.5 \pm 6.1 ^{ab}	127.5 \pm 78.4 ^c	100
78	<i>Sterilia</i> spp.	39.5 \pm 4.8 ^{ab}	28.9 \pm 9.2 ^a	50.1 \pm 16.5 ^{ab}	49.6 \pm 15.1 ^{ab}	64.5 \pm 20.1 ^b	50
135	<i>Sterilia</i> spp.	42.6 \pm 4.3 ^{ab}	43.5 \pm 2.8 ^{ab}	40.8 \pm 11.0 ^{ab}	45.3 \pm 5.0 ^{ab}	64.7 \pm 6.7 ^b	100
80	<i>Sterilia</i> spp.	27.4 \pm 18.8 ^a	36.7 \pm 4.1 ^{ab}	35.0 \pm 13.7 ^{ab}	40.2 \pm 18.4 ^{ab}	65.0 \pm 3.4 ^b	20–50
182	<i>Mucor</i> spp.	19.3 \pm 3.6 ^d	24.2 \pm 0.1 ^{ef}	23.2 \pm 1.6 ^e	10.9 \pm 11.5 ^b	4.02 \pm 2.4 ^a	50
166	<i>Mucor</i> spp.	19.4 \pm 1.3 ^d	24.5 \pm 0.5 ^{ef}	26.9 \pm 0.8 ^f	20.9 \pm 9.8 ^e	17.0 \pm 9.7 ^c	50–100
208	<i>Mucor</i> spp.	21.7 \pm 1.2 ^e	25.3 \pm 0.3 ^{ef}	22.2 \pm 0.1 ^e	13.5 \pm 9.3 ^{bc}	11.5 \pm 9.8 ^b	50
311	<i>Trichoderma</i> spp.	32.9 \pm 3.3 ^{ab}	17.3 \pm 0.4 ^a	26.4 \pm 0.2 ^a	30.9 \pm 1.6 ^{ab}	32.7 \pm 3.7 ^{ab}	>100
140	<i>Trichoderma</i> spp.	29.5 \pm 2.3 ^a	72.5 \pm 7.9 ^b	25.9 \pm 5.0 ^a	35.5 \pm 3.2 ^{ab}	46.6 \pm 2.4 ^{ab}	>100
1	<i>Penicillium</i> spp.	41.3 \pm 1.5 ^{jk}	32.6 \pm 5.6 ^g	34.8 \pm 2.4 ^{hi}	39.0 \pm 2.2 ^{ij}	50.9 \pm 5.1 ^{lm}	<5
5	<i>Penicillium</i> spp.	17.6 \pm 0.1 ^{ab}	29.8 \pm 3.4 ^{ef}	19.0 \pm 11.2 ^b	29.2 \pm 2.7 ^{ef}	36.6 \pm 4.7 ⁱ	<5
8	<i>Penicillium</i> spp.	19.3 \pm 11.8 ^{bc}	20.2 \pm 3.4 ^{bc}	30.9 \pm 11.2 ^{fg}	22.5 \pm 2.7 ^d	37.3 \pm 4.7 ⁱ	<5
44	<i>Penicillium</i> spp.	17.6 \pm 3.4 ^{ab}	24.3 \pm 0.6 ^{de}	16.5 \pm 5.9 ^{ab}	23.5 \pm 1.6 ^d	44.0 \pm 8.5 ^k	<5
6	<i>Penicillium</i> spp.	37.2 \pm 4.0 ⁱ	33.6 \pm 2.3 ^h	41.5 \pm 5.7 ^{jk}	48.5 \pm 11.1 ^l	58.0 \pm 10.8 ^m	<5
405	<i>Penicillium</i> spp.	17.3 \pm 8.4 ^{ab}	24.2 \pm 6.0 ^{de}	23.9 \pm 3.2 ^{de}	22.4 \pm 2.2 ^{cd}	36.4 \pm 5.2 ⁱ	<5
139	<i>Penicillium</i> spp.	17.4 \pm 12.9 ^{ab}	10.8 \pm 15.1 ^a	19.0 \pm 12.5 ^b	30.0 \pm 13.8 ^{ef}	>360 ⁿ	<5

Values represent the mean of four replicates. SD: standard deviation. Means from each genus with the same letter are not significantly different according to the LSD test ($p < 0.05$). (>) higher than. (<) lower than.

total incubation time, 360 h). Contrarily, *Mucor* spp. isolates showed the lowest values of this parameter with 100 mM of GP, especially isolate 182 remained only 4 h in the adaptation phase. For *Aspergillus* spp. and *Sterilia* spp., no significant differences in the length of the lag phase prior to growth with the increase of GP concentration was observed with respect to control. However, for isolates 128 (*Aspergillus* spp.), 133 and 80 (*Sterilia* spp.), the longest lag phases were also observed with 100 mM of GP ($p < 0.05$). For *Trichoderma* spp. the behavior

depended on the isolate tested. No significant differences between the lag phases in media supplied with GP and controls were observed for isolate 311, while for isolate 140, the longest lag phase was observed with 5 mM of GP. Likewise, for *Trichoderma* spp, for *Penicillium* spp. this growth parameter varied according to the isolates tested. In general, the longest lag phases were observed at the highest GP concentrations, and similar values were observed between controls and 5–50 mM media supplied with GP ($p < 0.05$).

With respect to EC50, the lowest value (<5 mM) was observed for *Penicillium* spp. While the highest value (>100 mM) was observed for *Trichoderma* isolates, followed by *Sterilia* spp. (isolates 133 and 135) with values of 100 mM, two *Aspergillus* spp. and one *Mucor* spp. (isolate 166) that had an EC50 value between 50 and 100 mM. In addition, EC50 values of 50 mM were found in *Sterilia* sp. (isolate 78) and *Mucor* spp. (isolates 182 and 208) and EC50 values of 20–50 mM in *Sterilia* spp. (isolate 80).

Figure 3 shows the growth rate of the 18 isolates in control media and in the presence of different GP concentrations at optimal water availability and temperature. The isolates developed with a different behavior in all GP concentrations assayed. The growth rates for *Sterilia* spp. and *Aspergillus* spp. decreased as GP increased, this reduction being more significant at the highest concentrations (Fig. 3A and D) ($p < 0.05$). For *Trichoderma* spp. the growth rate was constant with the increase of GP, e.g. isolate 140 showed a significant increase in growth rate with respect to the control with 50 mM (Fig. 3B). For all *Mucor* isolates, similar growth rate values were observed between the control media and the media supplied with 5 and 20 mM of GP. Furthermore, significant reductions of this parameter were found with 50 mM of GP (Fig. 3C). In general, for *Penicillium* spp., a decrease in growth rate was observed with the increase in GP concentration, being more evident with 50 mM. Only one isolate (139) did not develop at the highest herbicide concentration, while, with 5 and 20 mM of GP, similar or higher growth values were observed compared to the corresponding control (Fig. 3E) ($p < 0.05$). The isolates that showed the highest growth rates in media supplied with different GP concentrations were isolate 2 (*Aspergillus* spp.), isolates 182, 166 and 208 (*Mucor* spp.) and isolates 140 and 311 (*Trichoderma* spp.).

Discussion

The culturable mycota assay showed that the filamentous fungal count from the fields of Córdoba province varied according to the origin of the samples (exposure or non-exposure to pesticides) and the isolation media analyzed. Noticeable higher counts of culturable mycota in no-tillage or non-exposed soils to pesticides were observed compared to those in tillage and exposed soils both in controls as in the media supplied with GP. These results are partially in agreement with Cabello and Arambarri¹⁰ who evaluated the rhizospheric soil microfungi from a native forest (undisturbed and disturbed) from Buenos Aires, Argentina. They found highest counts in undisturbed soils. These authors informed that in these latter soils the number of colonies was 2-fold greater than the number found in disturbed forest at the same time. These results partially agree with the present study as the counts in non-exposed soils were higher than those in exposed soils; however, the differences were not as marked. The saprophytic mycota plays an important role in the cycles or nutrients in the ecosystems, because their capacity of fragmentation of organic polymeric matter allows the last mineralization by other microorganisms¹⁹. The high percentage of organic matter reported in soils without human intervention¹ can explain the high counts

observed in the non-exposed agricultural soil sampled in this work.

The most prevalent genera were *Fusarium* spp., *Aspergillus* spp., *Mucor* spp. and *Penicillium* spp., both in exposed and non-exposed soils. While *Trichoderma* spp., *Paecilomyces* spp. and *Verticillium* spp. were found only in exposed soils. These results agree partially with Carranza et al.¹⁵ and Nesci et al.³⁹, who informed a high frequency of *Aspergillus* spp., *Penicillium* spp., *Fusarium* spp., *Cladosporium* spp., *Trichoderma* spp. and *Paecilomyces* spp. isolated from maize fields in the south of Córdoba province exposed to pesticides and tillage practice. Only *Trichoderma* spp. and *Paecilomyces* spp. were the genera isolated only from non-exposed soils in the present study.

The main fungal genera isolated in this work were similar to a general isolation pattern found in non-rhizospheric soil ecosystems⁵ and in soils exposed to GP^{45,36}. The isolation of fungi such as *Penicillium*, *Aspergillus*, *Trichoderma* and *Cladosporium* is very common mainly in soil and organic plant material since the soil is the primary ecosystem for these fungi. They produce different propagation and resistance structures that provide them with adaptive features for environmental unfavorable conditions⁴² that explain their capacity for the fast invasion of the available substrate²³. These characteristics favor the maintenance of the species in the ecosystem. Wardle and Parkinson⁴⁸ and Reis Valpassos et al.⁴⁴ suggested that pesticides could change the saprophytic capacity of soil fungi or produce selective pressure that affects microbial activity favoring the development of more adapted genera. The presence of the same prevalent genera in exposed and non-exposed soils suggested that they possess major survival capacity to disturbers such as pesticide contamination and tillage.

Prevalent fungal genera counts in media supplied with GP showed differences according to the media analyzed. *Sterilia* spp. showed good development in all media, while *Mucor* spp. could only develop in media with GP as phosphorous source. The genus *Fusarium* spp. and *Penicillium* spp. developed in media with GP as nitrogen and phosphorous source, and *Aspergillus* spp. showed remarkable development with GP as phosphorous source. These results are indicating that the herbicide is used as a nutrient and the metabolic pathways vary according to the genus. The fungal enzymatic capacity and the rapid growth allow a fast invasion of the substrate to be used as a nutrient²⁷.

The length of the lag phase prior to growth and mycelial growth rate are the parameters used to indicate the effective development of the different isolates in the presence of GP at optimal a_w and temperature. The lag phase varied according to the isolate. The increase of GP concentration in the media produced an increase in the adaptation time in all isolates. However, this behavior was not observed in isolate 405 (*Mucor* spp.), as the lag phase values diminished significantly (80%) with 100 mM of GP with respect to control. These results agree with Barberis et al.⁷ who observed the same tendency in this parameter. They reported an increase in the lag phase at increasing GP concentrations (1–10 mM) in toxigenic *Aspergillus flavus* strains growing on maize-based medium.

The evaluation of this growth parameter suggests that the isolates needed in average an adaptation time higher than 30 h (*Aspergillus* spp., *Sterilia* spp., *Trichoderma* spp.




Figure 3 Growth rate of *Sterilia* spp. (133, 78, 135 and 80) (A), *Trichoderma* spp. (311 and 140) (B), *Mucor* spp. (182, 166 and 208) (C), *Aspergillus* spp. (128 and 2) (D), *Penicillium* spp. (1, 5, 8, 6, 44, 139 and 405) (E) in CZD media. a, i Mean values of growth rate with the same letter from each fungal genera are not statistically different according to the Fisher's LSD test ($p < 0.05$).

and *Mucor* spp.) and 24.7 h (*Penicillium* spp.) to start the exponential growth in media supplied with GP both at low and high concentrations. Although in the literature there are some studies regarding the impact of GP on fungal growth^{24,50,29}, there is scant information about the lag phase in relation to herbicides. In previous works, Caranza et al.¹⁵ reported lower lag phase values (18–23 h) than in the present study in non-toxicogenic *Aspergillus* section *Flavi* strains grown on low nutrient status media (soil based medium) conditioned at -0.70 of water potential and supplied with 5, 10 and 20 mM of GP. Contrarily for *Aspergillus* section *Nigri* strains, the lag phase values were higher (30–38 h) with 5, 10 and 20 mM of GP also in soil-based medium conditioned at -0.70 of water potential¹⁴. In other study, an increase in the lag phase at increasing GP concentrations (100–500 mM) was reported for *Aspergillus*

section *Flavi* strains grown on milled soybean extract agar¹². For *Fusarium verticillioides*, *Fusarium oxysporum* and *Fusarium graminearum* higher lag phases (36–90 h) were also reported in maize meal agar conditioned at 0.98 of a_w and supplied with 20, 30 and 50 mM of GP¹¹. Despite the fact that the evaluated isolates came from pesticide-exposed environments, they needed an acclimation period for their development in media with GP as a nutrient source. In this period, the hydrolytic enzymes are induced to metabolize the organophosphorus compounds. In the present study, the isolates showed variability in lag phase duration according to the GP concentration assayed, only one strain did not develop at the highest concentration evaluated.

Fungi are, in general, more tolerant to high xenobiotic concentration present in polluting sites than bacteria²². Results of EC50 showed that all isolates, except *Penicillium*

spp., were tolerant to GP. The *Trichoderma* isolates were the most tolerant (values >100 mM), followed by *Sterilia* spp. isolates 133 and 135 (values of 100 mM), two *Aspergillus* spp. and *Mucor* spp. isolate 166 with values between 50 and 100 mM. There is not information about this parameter for the estimation of sensitivity of fungal strains to GP. Previously, Nwachukwu and Osuji⁴⁰ reported tolerance to the herbicides atrazine, heptachlor and metolachlor of indigenous white rot fungus in soil extract agar. These fungi showed EC50 values of 0.011, 0.0038 mg/l and no inhibition or higher than 0.02 mg/l for atrazine, heptachlor and metolachlor, respectively. These EC50 values are notably lower than the ones found for *Trichoderma*, *Mucor*, *Aspergillus* and *Sterilia* isolates.

The growth rate varied according to the isolates tested. All of them could develop at all the GP concentrations assayed, except one *Penicillium* isolate (139) at the highest GP concentration. For *Penicillium* spp., *Sterilia* spp., and *Aspergillus* spp., the growth rates decreased with the increase of GP. This behavior was previously reported by Carranza et al.¹² with *Aspergillus oryzae* and non-toxicogenic *A. flavus* growing on milled soybean extract agar and supplied with higher concentrations of GP (100–500 mM). Contrarily, in other study, Carranza et al.¹⁵ reported a significant increase in the growth rate of *A. flavus* strains as the concentration of GP increased from 5 to 20 mM in soil extract solid medium conditioned at optimal water availability. Similar results were also found by Barberis et al.⁷ with toxigenic *Aspergillus* section *Flavi* strains grown in maize-based medium and supplied with 5 and 10 mM of GP. In the same way, a positive correlation between the GP degradation rate and the biomass of *A. oryzae* AF-02 was informed by XueHua et al.⁵⁰.

With respect to other genera, Krzysko-Lupicka and Orlik³² showed similar results with *Mucor*, *Penicillium* and *Trichoderma* strains isolated from soils. They grew exceptionally well in synthetic media supplied with GP as the sole source of phosphorus after a significant lag phase. While Klimek et al.²⁹ reported opposite results with a strain of *Penicillium crysogenum* as the concentrations of 5, 10 and 25 mM of GP in CZD medium (the herbicide replaced the nitrogen source) stimulated the fungal growth rate. Similar results were informed by Bujacz et al.⁹ who reported that this herbicide inhibited the *Penicillium notatum* (now *P. crysogenum*) growth in CZD medium with the highest GP concentrations, while with 1–0.5 mM the fungal growth significantly increased only in the media where the herbicide replaced the phosphorus source. With respect to *Mucor* spp., only one study performed *in situ* reported the effect of GP on this genus. Mandl et al.³⁶ reported the absence of *Mucor* spp. in the vineyard soil assay with GP. Similar growth results were also found in *Trichoderma viride* FRP3 isolated from Indonesian GP exposed soils⁴. Later, these authors reported the GP degradation capacity of this strain on soil and their promissory agriculture application³.

The biodegradation of GP, as for other organophosphorus compounds, depends on specific enzymatic and nonspecific enzymatic reactions. These processes occur *via* the secretion of phosphatase and other hydrolytic enzymes involved in the opening of phosphate bonds (P–O and P–S)²⁶. Genes encoding organophosphorus hydrolase and organophosphorus acid anhydrolase are expressed in several

microorganisms when they are exposed to organophosphorus compounds. This fact would induce the synthesis of the corresponding phosphatase. In addition, gene mutation and recombination are the most relevant genic processes that explain the microbial adaptation at contaminated sites¹⁷. This is consistent with several studies that have shown that the history of organophosphorus pesticide exposure would significantly improve the degradation rate^{28,25}. In the same way, it has been suggested that the most effective degradation process is related to functional microbial groups rather than to one microorganism⁴⁷.

Conclusion

The evaluation of all data showed that the two *Aspergillus* spp., two *Trichoderma* spp., three *Mucor* spp. and three *Sterilia* spp. (78, 133 and 135) isolates had the highest EC50 and growth rate values in media supplied with the herbicide. These isolates showed good growth performance after a variable acclimation period, indicating the inductive character of the GP degradation process. These results provide valuable data for further studies about the metabolic capacity of these promissory fungal species isolated from agricultural soils. These fungi could be potential candidates for enhancing GP removal both in pure and mixed cultures.

Funding

This work was financially supported by the Agencia Nacional de Promoción Científica y Tecnológica (ANPCYT-PICT-0943/14) and Secretaría de Ciencia y Técnica, Universidad Nacional de Río Cuarto (SECYT-UNRC-18/453).

Conflict of interest

The authors declare that they have no conflicts of interest.

Acknowledgments

This work was supported by the Agencia Nacional de Promoción Científica y Tecnológica (ANPCYT-PICT-0943/14), Secretaría de Ciencia y Técnica, Universidad Nacional de Río Cuarto (SECYT-UNRC-18/453) and Member of the Research Career of Consejo Nacional de Investigaciones Científicas y Técnicas (CONICET). The authors are specially grateful to Departamento de Microbiología e Inmunología of Universidad Nacional de Río Cuarto.

References

1. Allegrucci N, Cazau MC, Cabello MN, Arambarri AM. Análisis de las comunidades de microhongos de la hojarasca de *Scutia buxifolia* (*Rhamnaceae*) en el este de la Provincia de Buenos Aires, Argentina. *Darwiniana*. 2005;43:1–9.
2. Aparicio VC, De Gerónimo E, Marino D, Primost J, Carriquiriborde P, Costa JL. Environmental fate of glyphosate and aminomethylphosphonic acid in surface waters and soil of agricultural basins. *Chemosphere*. 2013;93:1866–73.
3. Arfarita N, Djuhari, Prasetya B, Imai T. The application of *Trichoderma viride* strain frp 3 for biodegradation of

- glyphosate herbicide in contaminated land. *Agrivita J Agric Sci.* 2016;38:275–81.
4. Arfarita N, Imai T, Kanno A, Higuchi T, Yamamoto K, Sekine M. Screening of soil-born fungi from forest soil using glyphosate herbicide as the sole source of phosphorus. *J Water Environ Technol.* 2011;9:391–400.
 5. Atlas RM, Bartha R. *Microbial ecology: fundamentals and applications.* 4th ed. The Singapore: Pearson Education Asia; 2000.
 6. Bansal OP. Degradation of pesticides. In: Rathore HS, Nollet LML, editors. *Pesticides, evaluation of environmental pollution.* London, New York: CRC Press, Taylor and Francis Group; 2012. p. 47–77.
 7. Barberis CL, Carranza CS, Chiacchiera SM, Magnoli CE. Influence of herbicide glyphosate on growth and aflatoxin B1 production by *Aspergillus* section *Flavi* strains isolated from soil on *in vitro* assay. *J Environ Sci Health B.* 2013;48:1070–9.
 8. Benbrook CM. Trends in glyphosate herbicide use in the United States and globally. *Environ Sci Eur.* 2016;28:1–15.
 9. Bujacz B, Wiczorek P, Krzysko-Lupicka T, Golab Z, Lejczak B, Kavfarski P. Organophosphonate utilization by the wild-type strain of *Penicillium notatum*. *Appl Environ Microbiol.* 1995;61:2905–10.
 10. Cabello M, Arambarri A. Diversity in soil fungi from undisturbed and disturbed *Celtis tala* and *Scutia buxifolia* forests in the eastern Buenos Aires province (Argentina). *Microbiol Res.* 2002;157:115–25.
 11. Carranza CS, Aluffi ME, Benito N, Magnoli K, Barberis CL, Magnoli CE. Effect of *in vitro* glyphosate on *Fusarium* spp. growth and disease severity in maize. *J Sci Food Agric.* 2019;99:5064–72.
 12. Carranza CS, Barberis CL, Chiacchiera SM, Dalcerro AM, Magnoli CE. Isolation of culturable mycobiota from agricultural soils and determination of tolerance to glyphosate of nontoxigenic *Aspergillus* section *Flavi* strains. *J Environ Sci Health B.* 2016;51:35–43.
 13. Carranza CS, Barberis CL, Chiacchiera SM, Magnoli CE. Assessment of growth of *Aspergillus* spp. from agricultural soils in the presence of glyphosate. *Rev Argent Microbiol.* 2017;49:384–93.
 14. Carranza CS, Barberis CL, Chiacchiera SM, Magnoli CE. Influence of the pesticides glyphosate, chlorpyrifos and atrazine on growth parameters of nonochratoxigenic *Aspergillus* section *Nigri* strains isolated from agricultural soils. *J Environ Sci Health B.* 2014;49:747–55.
 15. Carranza CS, Bergesio MV, Barberis CL, Chiacchiera SM, Magnoli CE. Survey of *Aspergillus* section *Flavi* presence in agricultural soils and effect of glyphosate on nontoxigenic *A. flavus* growth on soil-based medium. *J Appl Microbiol.* 2014;116:1229–40.
 16. Carranza CS, Regñicoli JP, Aluffi ME, Benito N, Chiacchiera SM, Barberis CL, Magnoli CE. Glyphosate *in vitro* removal and tolerance by *Aspergillus oryzae* in soil microcosms. *Int J Environ Sci Technol.* 2019, <http://dx.doi.org/10.1007/s13762-019-02347-x>.
 17. Castle LA, Siehl DL, Gorton R, Patten PA, Chen YH, Bertain S, Cho HJ, Duck N, Wong J, Liu D, Lassner MW. Discovery and directed evolution of a glyphosate tolerance gene. *Science.* 2004;304:1151–4.
 18. Castro Berman M, Marino DJG, Quiroga MV, Zagarese H. Occurrence and levels of glyphosate and AMPA in shallow lakes from the Pampean and Patagonian regions of Argentina. *Chemosphere.* 2018;200:513–22.
 19. Christensen M. A view of fungal ecology. *Mycologia.* 1989;81:1–19.
 20. DiRenzo A, Casanoves F, Balzarini G, Gonzalez L, Tablada M, Robledo C. Infostat. FCA. Argentina: Universidad Nacional de Córdoba; 2017 <http://www.infostat.com.ar>
 21. FAO Project concept paper. HEALTH: health in ecological agricultural learning. Prepared by the FAO programme for community IPM in Asia. Food and Agricultural; 2000.
 22. Fragoeiro S, Magan N. Impact of *Trametes versicolor* and *Phanerochaete chrysosporium* on differential breakdown of pesticide mixtures in soil microcosms at two water potentials and associated respiration and enzyme activity. *Int Biodeterior Biodegrad.* 2008;62:376–83.
 23. Frankland JC. Mechanisms in fungal successions. In: Wicklow DT, Carroll GC, editors. *The fungal community, its organization and role in the ecosystem.* New York: Marcel Dekker, Inc.; 1981. p. 403–26.
 24. Hasan HAH. Fungal utilization of organophosphate pesticides and their degradation by *Aspergillus flavus* and *A. sydowii* in soil. *Folia Microbiol (Praha).* 1999;44:77–84.
 25. Hernandez Guijarro K, Aparicio V, De Gerónimo E, Castellote M, Figuerola EL, Costa JL, Erijman L. Soil microbial communities and glyphosate decay in soils with different herbicide application history. *Sci Total Environ.* 2018;634:974–82.
 26. Huang X, He J, Yan X, Hong Q, Chen K, He Q, Zhang L, Liu X, Chuang S, Li S, Jiang J. Microbial catabolism of chemical herbicides: microbial resources, metabolic pathways and catabolic genes. *Pestic Biochem Physiol.* 2017;143:272–97.
 27. Jain R, Garg V. Enzymatic degradation of monocrotophohs by extracellular fungal OP ydrolases. *Appl Biochem Biotechnol.* 2013;171:1473–86.
 28. Karpuzcu ME, Sedlak DL, Stringfellow WT. Biotransformation of chlorpyrifos in riparian wetlands in agricultural watersheds: implications for wetland management. *J Hazard Mater.* 2013;244–5:111–20.
 29. Klimek M, Lejczak B, Kafarski P, Forlani G. Metabolism of the phosphonate herbicide glyphosate by a non-nitrate-utilizing strain of *Penicillium chrysogenum*. *Pest Manag Sci.* 2001;57:815–21.
 30. Koning LA, de Mol F, Gerowitt B. Effects of management by glyphosate or tillage on the weed vegetation in a field experiment. *Soil Tillage Res.* 2019;186:79–86.
 31. Krutz LJ, Shaner DL, Zablotowicz RM. Enhanced degradation and soil depth effects on the fate of atrazine and major metabolites in Colorado and Mississippi soils. *J Environ Qual.* 2010;39:1369–77.
 32. Krzysko-Lupicka T, Orlik A. Use of glyphosate as the sole source of phosphorus or carbon for the selection of soil-borne fungal strains capable to degrade this herbicide. *Chemosphere.* 1997;34:2601–5.
 33. Krzysko-Lupicka T, Sudol T. Interactions between glyphosate and autochthonous soil fungi surviving in aqueous solution of glyphosate. *Chemosphere.* 2008;71:1386–91.
 34. Lescano MR, Pizzul L, Castillo MDP, Zalazar CS. Glyphosate and aminomethylphosphonic acid degradation in biomixtures based on alfalfa straw, wheat stubble and river waste. *J Environ Manag.* 2018;228:451–7.
 35. Lupi L, Miglioranza KSB, Aparicio VC, Marino D, Bedmar F, Wunderlin DA. Occurrence of glyphosate and AMPA in an agricultural watershed from the southeastern region of Argentina. *Sci Total Environ.* 2015;536:687–94.
 36. Mandl K, Cantelmo C, Gruber E, Faber F, Friedrich B, Zaller JG. Effects of glyphosate, glufosinate and flazasulfuron based herbicides on Soil microorganisms in a vineyard. *Bull Environ Contam Toxicol.* 2018;101:562–9.
 37. Maqbool Z, Hussain S, Imran M, Mahmood F, Shahzad T, Ahmed Z, Azeem F, Muzammil S. Perspectives of using fungi as bioresource for bioremediation of pesticides in the environment: a critical review. *Environ Sci Pollut Res Int.* 2016;23:16904–25.
 38. Muzio J, Hilbert JA, Donato L, Arena P, Allende D. Technical comments data biodiesel from soybeans directorate general for energy and transport. Instituto Nacional de Tecnología Agropecuaria (INTA), document number IIR-BC-INF-14-08; 2008.
 39. Nesci A, Barros G, Castillo C, Etcheverry M. Soil fungal population in preharvest maize ecosystem in different tillage practices in Argentina. *Soil Tillage Res.* 2006;91:143–9.

40. Nwachukwu EO, Osuji JO. Bioremedial degradation of some herbicides by indigenous white rot fungus, *Lentinus subnudus*. *J Plant Sci*. 2007;2:619–24.
41. Okada E, Pérez D, De Gerónimo E, Aparicio V, Massone H, Costa JL. Non-point source pollution of glyphosate and AMPA in a rural basin from the southeast Pampas, Argentina. *Environ Sci Pollut Res Int*. 2018;25:15120–32.
42. Pitt JI, Hocking AD. *Fungi and food spoilage*. Dordrecht, Heidelberg, London, New York, Australia: Springer; 2009.
43. Primost JE, Marino DJG, Aparicio VC, Costa JL, Carriquiriborde P. Glyphosate and AMPA, “pseudo-persistent” pollutants under real-world agricultural management practices in the Mesopotamic Pampas agroecosystem, Argentina. *Environ Pollut*. 2017;229:771–9.
44. Reis Valpassos MA, Silva Cavalcante EG, Rodrigues Cassiolato AM, Alves MC. Effects of soil management systems on soil microbial activity, bulk density and chemical properties. *Pesq Agropec Bras*. 2001;36:1539–45.
45. Sailaja KK, Satyaprasad K. Degradation of glyphosate in soil and its effect on fungal population. *J Environ Sci Eng*. 2006;48:189–90.
46. Samson R, Houbraken J, Thrane U, Frisvad JC, Andersen B. *Food and indoor fungi*. CBS Laboratory Manual Series, vol. 2. Utrecht: CBS-KNAW Fungal Biodiversity Centre; 2010.
47. Wang B, Wu C, Liu W, Teng Y, Luo Y, Christie P, Guo D. Levels and patterns of organochlorine pesticides in agricultural soils in an area of extensive historical cotton cultivation in Henan province, China. *Environ Sci Pollut Res Int*. 2016;23:6680–9.
48. Wardle DA, Parkinson D. The influence of the herbicide glyphosate on interspecific interactions between four soil fungal species. *Mycol Res*. 1992;96:180–6.
49. Watanabe T. *Pictorial atlas of soil and seed fungi*. London, New York: CRC Press, Taylor and Francis Group; 2010.
50. XueHua W, GuiMing F, Yin W, DeBin G, YongHui C, YangFan L, XiaoFang W. Isolation and identification of glyphosate-degraded strain *Aspergillus oryzae* sp. A-F02 and its degradation characteristics. *Plant Dis Pests*. 2010;1:54–7.